

From: Kaushal, Sumesh
Sent: Monday, November 04, 2002 2:34 PM
To: STIC-Biotech/ChemLib
Subject: 09/755633: Interference and Sequence/Oligo Search

09/755633: Interference and Sequence/Oligo Search

Title: CANINE AND FELINE IMMUNOREGULATORY PROTEINS, NUCLEIC ACID MOLECULES, AND USES THEREOF
Inventor: YANG, SHUMIN

- SEQ ID NO: 4, 6, 7, 8, 9, 11, 18 and 19
- + Oligo search for SEQ ID NO: 4, 6, 7, 8, 9, 11, 18 and 19
- SEQ ID NO: 5 and 10 (PRT).

Prioriy Date: 5/29/1998

Thanks

S. Kaushal

CM1 12A07 AU1636

Ph: 703-305-6838

Mail Box: 11E12

GRF

Searcher: _____

Date Completed: _____

Searcher Prep/Review: _____

Clerical: _____

Online time: _____

TYPE OF SEARCH:

NA Sequences: 10

AA Sequences: _____

Protein Sequences: _____

Patent Literature: _____

Investigation: _____

Full text: _____

Patent Family: _____

Other: _____

VENDOR/COST (where applic.)


STN: _____

DIALOG: _____

Sequence Sys.: 1

WWW/Internet: _____

Other (specify): _____



Entrez PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed for **cannie IL-5** **Preview** **Go** **Clear**

Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
Utilities

PubMed Services
Journals Database
MESH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
ToxNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

- Search History will be lost after one hour of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Search	Most Recent Queries	Time	Result
#68 Related Articles for PubMed (Select 11440633)		14:51:07	425
#1 Search IL-5		14:40:19	5786
#49 Related Articles for PubMed (Select 7761472)		14:32:08	123
#48 Search Tavernier J[au]		14:31:21	235
#47 Search TF-1 AND IL-5 Field: All Fields		14:27:17	36
#46 Search TF-1 AND IL-5 Field: Pagination		14:27:08	0
#45 Search TF-1 AND IL_5 Field: Pagination		14:26:59	0
#44 Related Articles for PubMed (Select 8097324)		14:26:37	176
#42 Search #40 AND 3735 Field: Pagination		14:25:48	1
#40 Search McKenzie Field: Author, Limits: Publication Date from 1993 to 1993		14:24:56	148
#33 Search IL-5 AND TF-1 Field: All Fields		14:00:57	36
#37 Related Articles for PubMed (Select 9325013)		14:00:40	281
#32 Search IL-5 AND TF-1 Field: All Fields, Limits: Review		13:58:29	0
#29 Related Articles for PubMed (Select 9209411)		13:58:16	234
#30 Related Articles for PubMed (Select 9766809)		13:58:14	428
#15 Search IL-5 Field: Title, Limits: Review		13:55:19	40
#14 Search IL-5 Field: Title		13:51:00	583
#13 Search Interleukin-5		13:50:36	1
#12 Search #8 AND Yang S		13:50:16	0
#11 Search Weber ER AND Yang S		13:50:05	0
#10 Search Weber ER AND Yang S[Author]		13:49:50	0
#9 Search Weber ER AND Yang S[Author] Field: Author		13:49:42	0
#8 Search Weber ER Field: Author		13:49:11	46
#4 Related Articles for PubMed (Select 12113801)		13:48:53	99
#2 Search IL-5 Field: All Fields, Limits: Review		13:46:49	602

[Clear History](#)

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[PMC](#)[Taxonomy](#)[OMIM](#)[Books](#)Search **Nucleotide**for **cannie IL-5**[Preview](#)[Go](#)[Clear](#)[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)[About Entrez](#)

Search for Genes

LocusLink provides curated information for human, fruit fly, mouse, rat, and zebrafish.

Entrez Nucleotide

[Help](#) | [FAQ](#)

Batch Entrez: Upload a file of GI or accession numbers to retrieve sequences.

Check sequence revision history.

How to create VAMW links to Entrez.

[LinkOut](#)[Cubby](#)

Related resources

[ELAST](#)

Reference sequence project.

[Submit to GenBank](#)

- Search History will be lost after one hour of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Search


Most Recent Queries

#71 **Nucleotide Links for PubMed (Select 11440633)**#55 Search **cannie IL-5**#64 Search **cannie AND IL-5**#62 Search **txid9615[Organism]**#54 Search **IL-5**#53 Search **IL_5****Time Result**14:51:25 214:48:14 115814:47:32 014:45:46 2412014:40:53 115814:40:04 1158[Clear History](#)

Revised: July 5, 2002.

Disclaimer | Write to the Help Desk
NCBI | NLM | NIH

Mar 6, 2003 10:50:17



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

Utilities

PubMed Services

Journal Database

NCBI Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TriXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: J Interferon Cytokine Res 2000 Sep;20(9):779-85

Related Articles, Links

Canine interleukin-13: molecular cloning of full-length cDNA and expression of biologically active recombinant protein.

Yang S, Boroughs KL, McDermott MJ.

Heska Corporation, Fort Collins, CO 80525, USA. shuminyang@maxygen.com

Interleukin-13 (IL-13) regulates immune responses mediated by type 2 T helper lymphocytes (Th2) in the human and mouse. To study the function of this cytokine in the dog, we have isolated a cDNA that encodes the full-length canine IL-13 (CaIL-13) precursor polypeptide of 131 amino acids. CaIL-13 shares significant homology with the IL-13 amino acid sequences of cattle (54.1%), mouse (39.6%), and rat (36.6%) but shares the highest identity with human IL-13 (HuIL-13) (61.8%). The predicted CaIL-13 mature polypeptide of 111 residues was expressed in bacteria, and recombinant CaIL-13 (rCaIL-13) was isolated from inclusion bodies and refolded. rCaIL-13 stimulated the proliferation of TF-1 cells, which are derived from human erythroleukemia cells and respond to IL-13 as well as to a number of other human and murine cytokines. CaIL-13 mRNA was readily detectable by reverse transcriptase-polymerase chain reaction (RT-PCR) in cells from lymph nodes and peripheral blood. The gene sequence and biologically active recombinant protein for CaIL-13 will be useful reagents to determine the role of IL-13 in the regulation of canine immune responses.

PMID: 11032397 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

May 6, 2003 10:50:17



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed for Go Clear

☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed

Overview

Help: A-Z

Tutorial

New/Noteworthy

Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Pubby

Related Resources

Order Documents

NLM Gateway

ToxNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: Growth Factors 1993;8(2):87-97

Related Articles Links

A model for the interaction of the GM-CSF, IL-3 and IL-5 receptors with their ligands.

Goodall GJ, Bagley CJ, Vadas MA, Lopez AF.

Hanson Centre for Cancer Research, Division of Human Immunology, Institute of Medical and Veterinary Science, Adelaide, South Australia.

The high affinity receptors for GM-CSF, IL-3 and IL-5 are heterodimers consisting of a ligand-specific alpha chain and a common beta chain. These proteins are members of a family of proteins known as the "cytokine receptor family" which is characterized by the presence of a 200-residue ligand-binding module. The GM-CSF, IL-3 and IL-5 receptor alpha chains constitute a distinct subgroup and share features not found in other members of the cytokine receptor family, features which we propose to be important for their interaction with the common beta chain and for their binding of the structurally-related ligands. The growth hormone receptor is a well-characterized member of the cytokine receptor family. Based on the structure of the complex between growth hormone and its receptor, we have proposed sites of contact between the GM-CSF, IL-3 and IL-5 receptors and their cognate ligands.

Publication Types:


- Review
- Review, Tutorial

PMID: 8466757 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

Mar 6, 2003 10:50:17



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed for Go Clear

☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
Statistics

PubMed Services
Journals Database
MESH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

1: Exp Hematol 2000 Mar;28(3):231-43

Related Articles, Links

PubMed Reference
FULL-TEXT ARTICLE

A model for assembly and activation of the GM-CSF, IL-3 and IL-5 receptors: insights from activated mutants of the common beta subunit.

D'Andrea RJ, Gonda TJ.

Hanson Centre for Cancer Research and, Adelaide, South Australia, Australia.

Granulocyte-macrophage colony stimulating factor (GM-CSF), Interleukin-3 (IL-3) and Interleukin-5 (IL-5) have overlapping, pleiotropic effects on hematopoietic cells, including neutrophils, eosinophils, monocytes and early progenitor cells. The high-affinity receptors for human GM-CSF, IL-3, and IL-5 share a common beta-subunit (hbeta(c)), which is essential for signalling and plays a major role in recruiting intracellular signalling molecules. While activation of the cytoplasmic tyrosine kinase JAK2 appears to be the initiating event for signalling, the immediate events that trigger this are still unclear. We have isolated a number of activated mutants of hbeta(c), which can be grouped into classes defined by their state of receptor phosphorylation, their requirement for alpha subunit as a cofactor, and their activities in primary cells and cell lines. We discuss these findings with regard to the stoichiometry, activation, and signalling of the normal GM-CSF/IL-3/IL-5 receptor complexes. Specifically, this work has implications for the role of the ligand-specific alpha-subunits in initiating the signalling through the beta-subunit, the role of beta subunit dimerization as a receptor trigger, and the function of receptor tyrosine phosphorylation in generating growth and survival signals. Based on the properties of the activated mutants and the recent structures of erythropoietin receptor (Epo-R) complexes, we propose a model in which (1) activation of hbeta(c) can occur via alternative states that differ with respect to stoichiometry and subunit assembly, but which all mediate proliferative responses, and (2) each of the different classes of activated mutants mimics one of these alternative states.


Publication Types:

- Review
- Review, Tutorial

PMID: 10720688 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed for Go Clear

☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed

Overview

Help: FAQ

Tutorial

New/Noteworthy

Books

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Under Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts


Clinical Trials.gov

PubMed Central

Privacy Policy

1: Expert Opin Investig Drugs 2000 Mar;9(3):491-6

Related Articles, Links



IL-5: biology and potential therapeutic applications.

Weltman JK, Karim AS.

Department of Medicine, Brown University School of Medicine, Providence, RI 02912, USA.
joel.weltman@brown.edu

IL-5 is the predominant cytokine associated with antigen-induced eosinophilic inflammation in the lung. The activation of Th-2 cells leads to the production of IL-5. The pro-eosinophilic effects of IL-5 include: (1) enhanced replication and differentiation of eosinophilic myelocytes; (2) enhanced degranulation of eosinophils; (3) prolonged survival time of eosinophils; and (4) enhanced adhesion of eosinophils. The effects of IL-5 are mediated via the interaction of IL-5 with receptors (IL-5R) that are expressed on the eosinophil cell membrane. Intracellular signalling produced by occupation of the IL-5R by IL-5 occurs via the JAK-STAT system. IL-5 is a 45 kDa glycoprotein consisting of two identical polypeptide chains. The 5'-promoter region of the IL-5 gene contains elements that are down-regulated by glucocorticoids. Anti-IL-5 reagents have the potential to suppress IL-5 activity without the side effects of glucocorticoids. Studies using monoclonal antibodies (mAbs) against IL-5 have established the feasibility of suppressing eosinophilic inflammation by specifically blocking IL-5 activity. Studies with antisense IL-5 are beginning to provide the basis for non-glucocorticoid, sequence-specific oligonucleotide inhibitors of IL-5. Research has begun on the development of mAbs and antisense oligonucleotide inhibitors of IL-5 that can be inhaled and applied topically.

Publication Types:

- Review
- Review, Tutorial

PMID: 11060690 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text


Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

NCBI 2000-01-17



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed for Go Clear

☒ Limits Preview/Index History Clipboard Details

About Entrez

Help

Text Version

Entrez PubMed

Overview

Help FA2

Tutorial

New! Noteworthy

Utilities

PubMed Services

Journals Database

Medline Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Caddy

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Nippon Rinsho 2001 Oct;59(10):1894-9

Related Articles. Links

[Inflammatory cytokines (IL-4, IL-5 and IL-13)]

[Article in Japanese]

Maeda S, Yanagihara Y.

Clinical Research Center, National Sagami Hospital.

The polarized Th2 cells play an important role in the pathogenesis of atopic asthma as well as in the induction of airway inflammation. Th2 cytokines, such as IL-4, IL-5 and IL-13, are pivotal in regulating the allergic phenotype, the IgE response or the inflammatory cell-mediated function. Selective inhibition of Th2 cytokines by pharmacologic agents, including anti-cytokine blocking antibody, cytokine mutant and soluble cytokine receptor, will contribute to asthma therapy. Strategies based on blocking key signaling cytokines are also discussed.

Publication Types:

- Review
- Review, Tutorial

PMID: 11676128 [PubMed - indexed for MEDLINE]


Display Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health & Human Services

Freedom of Information Act | Disclaimer



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed for Go Clear

☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
Utilities

PubMed Services

Journals Database
Mesh Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

☐ 1: Pharmacol Ther 2002 Jun;94(3):253-64

Related Articles Links

FULL-TEXT ARTICLE

Interleukins-4, -5, and -13: emerging therapeutic targets in allergic disease.

Foster PS, Martinez-Moczygemba M, Huston DP, Corry DB.

Division of Biochemistry and Molecular Biology, John Curtin School of Medical Research, Australian National University, Canberra, ACT 0200 Australia.

For the first time, allergic diseases have emerged as major public health concerns. Highly effective therapies for allergic disease now exist, but are plagued by serious side effects and the fact that a significant minority of patients remains unresponsive. Studies from many laboratories have established that T helper type 2 (T(H)2) cytokines contribute importantly to diseases such as asthma, and therapeutic strategies that target the key T(H)2 cytokines are of potential benefit in allergic disease. In this article, we will review the biology of the T(H)2 cytokines interleukin (IL)-4, IL-5, and IL-13 and their receptors, and will consider several novel strategies to neutralize these molecules in human and experimental asthma. While promising, newer therapies face a gauntlet of developmental challenges, but offer the hope of reducing allergic diseases once again to minor public health concerns. Copyright 2002 Elsevier Science Inc.

Publication Types:


- Review
- Review, Tutorial

PMID: 12113801 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

Mar 6, 2003 10:50:17



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show: 20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help [FAQ]

Tutorial

New/Noteworthy

Errata

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: J Interferon Cytokine Res 2001 Jun;21(6):361-7

Related Articles, Links

Canine interleukin-5: molecular characterization of the gene and expression of biologically active recombinant protein.

Yang S, Sellins KS, Weber E, McCall C.

Heska Corporation, Fort Collins, CO 80525, USA.

Interleukin-5 (IL-5), which is produced primarily by type 2 T helper lymphocytes (Th2), is an eosinophil differentiation and activation factor. Increased numbers of eosinophils in peripheral blood or tissues (eosinophilia) are observed in asthmatic human patients, in animals with helminth infections, and in dogs with allergic diseases. Antagonism of IL-5 activity is being explored as a potential treatment of a number of disease conditions associated with eosinophils in animal models. In order to study the expression and function of this cytokine in the dog, we have isolated and characterized the canine IL-5 gene. The canine IL-5 polypeptide deduced from the cDNA is composed of 134 amino acids that share varying degrees of homology with IL-5 isolated from several mammals. The genomic structure of the canine IL-5 gene consists of four exons and three introns in the coding region, similar to that of the previously characterized human and mouse IL-5 genes. Recombinant canine IL-5 protein, expressed in *Pichia pastoris*, is biologically active in a cell proliferation assay. Canine IL-5 gene sequences and the biologically active protein described in this study will be useful reagents for future studies of this cytokine in physiologic processes and in pathologic conditions of the dog.

PMID: 11440633 [PubMed - indexed for MEDLINE]

Display

Abstract

Show: 20

Sort

Send to

Text


Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

NCBI.nlm.nih.gov



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search PubMed

for

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show: 20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help [A-Z]

Tutorial

New/Noteworthy

Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: EMBO J 1994 Nov 1;13(21):5176-85

Related Articles, Links

Three residues in the common beta chain of the human GM-CSF, IL-3 and IL-5 receptors are essential for GM-CSF and IL-5 but not IL-3 high affinity binding and interact with Glu21 of GM-CSF.

Woodcock JM, Zacharakis B, Plaetinck G, Bagley CJ, Qiyu S, Hercus TR, Tavernier J, Lopez AF.

Division of Human Immunology, Hanson Centre for Cancer Research, Institute of Medical and Veterinary Science, Adelaide, South Australia.

The beta subunit (beta c) of the receptors for human granulocyte macrophage colony stimulating factor (GM-CSF), interleukin-3 (IL-3) and interleukin-5 (IL-5) is essential for high affinity ligand-binding and signal transduction. An important feature of this subunit is its common nature, being able to interact with GM-CSF, IL-3 and IL-5. Analogous common subunits have also been identified in other receptor systems including gp130 and the IL-2 receptor gamma subunit. It is not clear how common receptor subunits bind multiple ligands. We have used site-directed mutagenesis and binding assays with radiolabelled GM-CSF, IL-3 and IL-5 to identify residues in the beta c subunit involved in affinity conversion for each ligand. Alanine substitutions in the region Tyr365-Ile368 in beta c showed that Tyr365, His367 and Ile368 were required for GM-CSF and IL-5 high affinity binding, whereas Glu366 was unimportant. In contrast, alanine substitutions of these residues only marginally reduced the conversion of IL-3 binding to high affinity by beta c. To identify likely contact points in GM-CSF involved in binding to the 365-368 beta c region we used the GM-CSF mutant eco E21R which is unable to interact with wild-type beta c whilst retaining full GM-CSF receptor alpha chain binding. Eco E21R exhibited greater binding affinity to receptor alpha beta complexes composed of mutant beta chains Y365A, H367A and I368A than to those composed of wild-type beta c or mutant E366A. These results (i) identify the residues Tyr365, His367 and Ile368 as critical for affinity conversion by beta c, (ii) show that high affinity binding of GM-CSF and IL-5 can be dissociated from IL-3 and (iii) suggest that Tyr365, His367 and Ile368 in beta c interact with Glu21 of GM-CSF.

PMID: 7957082 [PubMed - indexed for MEDLINE]

Display

Abstract

Show: 20

Sort

Send to


Text

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Books

Search

Limits

Preview/Index

History

Clipboard

Details

Display

Show:

Send to ☐ 1: AF331920. Canis familiaris ...[gi:15919182]

Links

LOCUS AF331920 1658 bp DNA linear MAM 04-OCT-2001

DEFINITION Canis familiaris interleukin-5 gene, complete cds.

ACCESSION AF331920

VERSION AF331920.1 GI:15919182

KEYWORDS .

SOURCE Canis familiaris (dog)

ORGANISM Canis familiarisEukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Carnivora; Fissipedia; Canidae; Canis.

REFERENCE 1 (bases 1 to 1658)

AUTHORS Yang, S., Sellins, K.S., Weber, E. and McCall, C.

TITLE Canine interleukin-5: molecular characterization of the gene and
expression of biologically active recombinant protein

JOURNAL J. Interferon Cytokine Res. 21 (6), 361-367 (2001)

MEDLINE 11334408PubMed 11440638

REFERENCE 2 (bases 1 to 1658)

AUTHORS Yang, S.

TITLE Direct Submission

JOURNAL Submitted (22-FEC-2000) Immunology, Heska Corporation, 1613
Prospect Parkway, Ft Collins, CO 80525, USA

FEATURES

source Location/Qualifiers

1..1658

/organism="Canis familiaris"

/db_xref="taxon:9615"

mRNA /join(1..170,374..406,1276..1404,1523..1658)

/product="interleukin-5"

5'UTR 1..26

CDS /join(27..170,374..406,1276..1404,1523..1621)

/note="IL-5"

/codon_start=1

/product="interleukin-5"

/protein_id="AAL10716.1"

/db_xref="GI:15919183"

/translation="MRMLNLNLSTLALGAAYVSGAVENHNNPLVAETLTLLSTHRTWL
LUNNNLHPTPENKXHQDTHVFDGHTLKH,TAHSHAVTSLFQNLSLIKHIERQF
KRCAGELWRVTFELDYIQVFLAVINTEWTLER"

3'UTR 1..2..165:

BASE COUNT 839 a 257 g 363 g 162 t

ORIGIN

```

1 aggcacacac tgaacattc agagctatga gaatgcttct gaatttgagt ttgctagctc
61 ttggggctgc ctatgttct gctttrgtg tagaaaaatc catgaataga ctggtggcag
121 agaccttgac actgctctcc actcatcgaa cttggctgat aggcgatggg gtaattttct
181 ttttgattcc tacagtcttt aaaatgcacg gtaatttgtt ggtggtggct agtttttaaa
241 gatccattat caataatgaa gtaateagtr ttaataatct ataattgggt accatggtac
301 taagaaact tataataaaa gtatgaacc ttaataata ttaaaatga atgttggttc
361 attttttt taaagctga taattctac taaagaaat aaaaatgaa tttaaattat

```

1021 ctcgtaagca tttatttttc attaatcatt tcatttatat catttgtaac acttctcagt
 1081 aattatataa acatcattta cttatggtaa ttatagctta gtataagggtg gtttccacc
 1141 tggaaaagac acaagtaaaa acctcttggg agaagggaac ttgtgtaaac cccacaaaac
 1201 aaagtctaac tttttggacc aaatttttat gccttgtttt gatgaattat attttttaa
 1261 atcttccctc ttttagcacc actgtgcatt aaagaagttt ttcagggtat agacacattg
 1321 aagaacccaa ctgcccacgg ggaggctgtg gataaactat tccaaaactt gtcttcaata
 1381 aaagaacaca tagagcgcca aaaagtaagt taaagacatt tggcaaaaac ttaagtatat
 1441 ttgtctgaet ctgcctgttt tttttttttt tttttacaag aattgacagt ttctataat
 1501 atctccctctg ttcttttaac agaaaagggtg tgcaggagaa agatggagag tgacaaggtt
 1561 cctagactac ctgcaagtat ttcttgggtg aataaacacc gagtggacac cggaaagtgt
 1621 agaacaaaac ggcttattgt agtgggaagt ttgggaga

//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)
 NCBI | NLM | NIH

Mar 6 2003 14:06:04



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Books

Search

for

Limits

Preview/Index

History

Clipboard

Details

Display

Show:

Send to

1: AF331920. Canis familiaris ...[gi:15919182]

Links

LOCUS AF331920 468 bp DNA linear MAM 04-OCT-2001

DEFINITION Canis familiaris interleukin-5 gene, complete cds.

ACCESSION AF331920 REGION: join(1..170,374..406,1276..1404,1523..1658)

VERSION AF331920.1 GI:15919182

KEYWORDS

SOURCE Canis familiaris (dog)

ORGANISM Canis familiaris

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Carnivora; Fissipedia; Canidae; Canis.

REFERENCE 1 (bases 1 to 468)

AUTHORS Yang, S., Mellins, E.S., Weber, E. and McCall, C.

TITLE Canine interleukin-5: molecular characterization of the gene and expression of biologically active recombinant protein

JOURNAL J. Interferon Cytokine Res. 21 (6), 361-367 (2001)

MEDLINE 11364405

PubMed 14406331

REFERENCE 1 (bases 1 to 468)

AUTHORS Yang, S.

TITLE Direct Submission

JOURNAL Submitted (22-DEC-2000) Immunology, Heska Corporation, 1613 Prospect Parkway, Ft Collins, CO 80525, USA

FEATURES Location:Qualifiers

source

1..468

-organism="Canis familiaris"

-db_xref="taxon:9615"

mRNA

1..468

-product="interleukin-5"

5'UTR

1..26

CDS

17..431

-note="IL-5"

-codon_start=1

-product="interleukin-5"

-protein_id="AA110716.1"

-db_xref="GI:15919183"

-translation="MRMLLNLSLALGAAYVRAFAVENFKNPLVAETLTLLSTHPTWL

IGDENIMIPTFENANHQQLIKENFQQLTIRNQTANGEAVIKIPQLSLIKENIERQK

KRWAGFKWRVTEFLDYLOVFLGVINTEWTPES"

3'UTR

432..468

PAGE COUNT 165 a 165 b 165 c 165 d

ORIGIN

```

1 aggcacaaac tgaacatttc agagctatga gaatgcttct gaatttgagt ttgctagctc
61 ttggggcttc ctatgtttct gcttttgctg tagaaaatcc catgaataga ctggtggcag
121 agaccttgac actgctctcc actcatcgaa cttggctgat aggcatggg aacctgatga
181 ttctactcc tgaataataa aatcaccaac tgtgcattaa agaagtttt cagggtatag
241 acacattgaa aaacaaact gccacaggg aggetgtgga taaactattc cnaaacttat
301 atttaataa agacacata gacacacaa aaaaaattt tgcagtaaaa agatggagat
361 taaataaagt atgaataaac ctgaacat atcttttttt aataaataa aatgaacaaa

```

revised July 5, 2001

<http://www.ncbi.nlm.nih.gov/

RPS-BLAST 2.2.5 [Nov-16-2002]

Query= g|114191|g|114191|114191.1[6.1] aa (114) interleukin-5 [Canis familiaris]
(134 letters)

Database: cdd.v1.60
10,013 PSSMs; 2,494,783 total columns

Click on boxes for multiple alignments



Show Domain Relatives

- .. This CD alignment includes 3D structure. To display structure, download [Cn3D!](#)

PSSMs producing significant alignments:

Score
(bits) E
value

- gnlCDD8195 pfam02025, IL5, Interleukin 5

186 7e-49

- gnlCDD8195, pfam02025, IL5, Interleukin 5.

CD-Length = 108 residues, 99.1% aligned
Score = 186 bits (473), Expect = 7e-49

Query: 25 PMNRVASTLTLSFHTWLIQDGNLM:PTFENKNNHQLCIKEVFQGLDTLKNQTAHGEAV 84
Sbjct: 2 EMSALVKETLALSTHRTLLSNETLR:PVPTHKNNHQLCTEPIFQGLDTLKNQTAQGGAV 61

Query: 85 DKLPQNLSLINEHIERQKNRQAGERWVTFELDYLDQVFLGVINTEWT 131
Sbjct: 62 ETLPQNI SLIKKYIDRQKKKCGEERRKVKQFLDYLDQVFLGVINTEWT 108



CDART: Conserved Domain Architecture Retrieval Tool

[New Query](#)[Overview](#)[PubMed](#)[Nucleotide](#)[Protein](#)[Structure](#)[Taxonomy](#)[Help?](#)

0

100

IL5


Query


Similar domain architectures

21 Sequences

Therid

Interleukin-5 prec

Result page: [Previous](#) 1 [Next](#)**Subset** by Taxonomy**Subset** by selected domains:☐  pfam02025 Interleukin 5.[Help](#) | [Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

	BLAST	Protein	Structure	PubMed	Taxonomy
	Genome	Nucleotide	3D-Domains	Books	Help























Query: gi|15919183 interleukin-5 [Canis familiaris]
Matching gi: 15919181

[Best hits](#)
[Common Tree](#)
[Taxonomy Report](#)
[3D structures](#)
[CDD-Search](#)
[GI list](#)

22 BLAST hits to 18 unique species Sort by taxonomy proximity

☐ Archaea
☐ Bacteria
☒ Metazoa
☐ Fungi
☐ Plants
☐ Viruses
☐ Other Eukaryotae

Keep only Cut-Off [Select](#) [Reset](#)

134 aa	SCORE	E	ACCESSION	GI	PROTEIN DESCRIPTION
	591	23	AAC64506	3238519	interleukin-5 [Felis catus]
	586	23	AAC27516	3342342	interleukin 5 [Felis catus]
	573	31	AB18368	4461317	Interleukin 5 [Sus scrofa]
	571	36	AF056791	569073	interleukin-5; IL-5 [Canis familiaris]
	567	21	AAB01344	1243344	interleukin-5 [Equus caballus]
	551	21	CAA91704	1113141	interleukin-5 [Bos taurus]
	536	31	AAE10719	891772	interleukin-5 [Ovis aries]
	468	3	AAC05717	2961151	interleukin-5 [Felis catus]
	448	21	AAK43015	1537200	interleukin-5 [Saimiri sciureus]
	430	11	CAA13141	133133	B cell differentiation factor I [Homo sapi
	429	31	AAAG9411	514340	interleukin 5
	426	21	AAAG6710	644414	interleukin-5
	424	11	CAA13141	133133	B cell differentiation factor I [Homo sapi
	413	31	AAE01700	213113	interleukin-5 [Cavia porcellus]
	386	31	AAG16721	1034542	interleukin-5 [Sigmodon hispidus]
	386	21	AAK43015	1537200	interleukin 5
	374	31	CAAC9609	124334	IL-5 [Mus musculus]
	370	30	AAL14462	5004336	interleukin-5 [Macropus eugenii]
	359	31	CAAG283	311145	interleukin 5 [Rattus rattus]
	352	31	CAAG0587	4151563	interleukin-5 [Rattus norvegicus]
	349	21	CHULA	1311008	Chain A, Mol_id: 1; Molecule: Interleukin-
	108	20	AAD37461	5006324	interleukin-5 [Sminthopsis macroura]